



Membrane Potential and Glucose Homeostasis: Understanding the Role of Ion Channels in Diabetes Mellitus

***¹Ezeagwuna Chiamaka, ²Nnodim Johnkennedy, ¹Udegbulem Daniel Nwanneamaka, and ¹Nwodo Divine Akachukwu**

¹Department of Medical Laboratory Science, Madonna University, Elele Rivers State Nigeria.

²Department of Medical Laboratory Science, Imo State University, Owerri, Nigeria.

Corresponding Author: Ezeagwuna Chiamaka

Department of Medical Laboratory Science, Madonna University, Elele Rivers State Nigeria..

Received Date: 30 Aug. 2026

Published Date: 13 Sept. 2026

Abstract

Pancreatic β -cells, insulin release, peripheral tissues and cellular signalling pathways interact in a coordinated manner for the maintenance of glucose homeostasis. Membrane potential of excitable cells is a critical, although sometimes neglected, part of this process. The membrane potential is established by the uneven distribution of ions on the two sides of the plasma membrane and is mainly regulated by ion channels and transporters. In pancreatic β -cells, changes in membrane potential are generated through ATP-sensitive potassium (K_{ATP}) channels, in response to variations in glucose metabolism. Enhanced glucose metabolism causes increased intracellular ATP, leading in the closure of K_{ATP} -channels, membrane depolarisation, activation of voltage-dependent calcium (Ca^{2+}) channels and Ca^{2+} influx, which increases insulin secretion. Voltage-dependent potassium (K_V), calcium-activated potassium (K_{Ca}), calcium channels and other ion channels later modulate electrical activity and insulin release. These pathways can be disturbed in diabetes mellitus by prolonged hyperglycaemia, lipotoxicity, oxidative stress, mitochondrial dysfunction and altered intracellular calcium homeostasis. Thus, aberrant ion-channel activity may lead to poor β -cell electrical activity, decreased insulin production and increasing metabolic dysfunction. Ion channels are also involved in vascular and neuronal consequences of diabetes. Thus, understanding the membrane potential and modulation of ion channels provides a crucial foundation to comprehend the physiological basis of glucose dysregulation and may discover novel biomarkers and therapeutic targets in diabetes mellitus.

Keywords: Membrane potential; glucose homeostasis; ion channels; potassium channels; calcium channels; K_{ATP} channels; pancreatic β -cells; insulin secretion; diabetes mellitus; oxidative stress.

Introduction

Glucose homeostasis is the mechanism by which blood glucose levels are maintained within a healthy range despite variations in food consumption, physical activity and metabolic demand. This mechanism is highly dependent on the coordinated activity of insulin, glucagon, liver, skeletal muscle, adipose tissue and pancreatic islet cells. Among these components, pancreatic β -cells are central, sensing changes in circulating glucose and releasing insulin accordingly [1].

The electrical characteristics of the β -cells allow a quick response to glucose. Membrane potential: The electrical gradient generated by ion transport across the plasma membrane. In β -cells, this electrical state is regulated by glucose metabolism, coupling nutrient sensing to insulin production. Thus, ion channels are molecular translators of metabolic changes into electrical impulses and ultimately the production of hormones. K_{ATP} , voltage gated calcium and voltage gated potassium channels are particularly crucial in this mechanism [2].



Recent work has also emphasised the significance of membrane potential and calcium signalling in diabetes. Maduike, Nnodim and Nnodim considered the membrane potential as a possible laboratory biomarker in type 2 diabetes, and related it to ion channels, β -cell activity, oxidative stress and glycaemic control. Similarly, Nnodim and Ozoemena stated that calcium signalling and its modulation by ion channels and transporters are critical for insulin secretion and glucose homeostasis, and dysregulation of calcium homeostasis is associated with β -cell failure in diabetes[3].

Cell Function and Membrane Potential

Membrane potential is the charge differential across a cell membrane. This is due to the differential in ionic concentration across the plasma membrane and the selective permeability to ions like K^+ , Na^+ , Ca^{2+} and Cl^- . The electrical state of the cell is determined by these charged particles controlled by ion channels.

Potassium conductance plays a particularly essential role in setting the resting membrane potential of pancreatic β -cells. K_{ATP} channels are relatively open at low glucose levels, facilitating K^+ efflux and a negative membrane potential. This blocks electrical activity in the β -cell while reducing calcium entrance and insulin secretion. As glucose concentration rises, glucose enters the β -cell and undergoes glycolysis and mitochondrial metabolism. The consequent rise in ATP compared to ADP blocks K_{ATP} channels. The decreased potassium conductance induces a depolarisation of the membrane. This depolarisation brings the membrane potential closer to the activation threshold of voltage-dependent calcium[4] channels[4].

K ATP Channel and Glucose Sensing

The K_{ATP} channel is among the most critical linkages between electrical activity and cellular metabolism. In pancreatic β -cells, it is mostly constituted of Kir6.2 potassium-channel subunits and SUR1 regulatory subunits. Its activity is controlled by the quantities of intracellular ATP and ADP. At low glucose concentrations, the low ATP synthesis keeps K_{ATP} channels sufficiently active to maintain membrane hyperpolarisation. Metabolism increases the intracellular ATP/ADP ratio after a rise in glucose. The ATP-dependent blockage of K_{ATP} channels reduced K^+ outflow and depolarised the plasma membrane[5].

Thus, K_{ATP} channels are metabolic sensors and electrical switches coupling glucose metabolism to insulin production.

Calcium channels in insulin secretion

Insulin secretion is not merely a consequence of membrane depolarisation. It has to be translated into an intracellular calcium signal. The β -cell membrane depolarises to the appropriate threshold, voltage-dependent Ca^{2+} channels open, and Ca^{2+} enters the cell.

The subsequent rise of intracellular Ca^{2+} concentration activates the machinery for exocytosis of insulin-containing secretory granules. Thus, calcium is an important second messenger coupling electrical activity and hormone release[6]. Ca^{2+} signalling is also modulated by intracellular storage, calcium pumps, exchangers and other calcium-permeable channels. According to Nnodim and Ozoemena, the voltage-gated calcium channels, transient receptor potential channels, store-operated calcium-entry routes and calcium ATPases are essential regulators of calcium transport and intracellular calcium homeostasis in diabetes.

Thus, interruption at any stage in Ca^{2+} entry, storage, buffering, or extrusion can alter β -cell function. Excessive or prolonged increase of intracellular Ca^{2+} may also lead to cellular stress, mitochondrial malfunction and β -cell injury[7].

Potassium Channel Function in Membrane Repolarisation

Although the main players in glucose-induced depolarisation are the K_{ATP} channels, other potassium channels modulate the duration and pattern of beta-cell electrical activity. After depolarisation, voltage-dependent potassium channels open, allowing K^+ to exit from the cell. This leads to membrane repolarisation and prevents the duration of calcium-channel activation[8].

The calcium-potassium channel interaction is thus important in the rhythmic electrical activity of beta-cells. Potassium currents are activated by depolarisation and calcium entering to bring the membrane back to a more negative potential. This reduces calcium influx. This helps make insulin secretion pulsatile.

Disruption of this finely tuned balance might lead to improper membrane excitability and aberrant insulin secretion [10].

Dysfunction of Membrane Potential in Diabetes Mellitus

Diabetes mellitus is associated with severe metabolic alterations which can impact ion-channel function. Chronic hyperglycaemia promotes oxidative stress, changes mitochondrial metabolism and may interfere with intracellular signalling. Lipotoxicity, inflammation, endoplasmic-reticulum stress and mitochondrial malfunction may further impair β -cell function [11].

Ion channels are sensitive to metabolic and cellular circumstances and perturbations can change channel expression, activity or regulation. Abnormal K_{ATP}-channel activity might disrupt the natural link between glucose metabolism and membrane depolarisation. In a similar way, abnormal voltage-dependent calcium-channel activity may lead to insufficient or excessive calcium influx[12]. At the level of calcium signalling, disturbance of intracellular calcium control may lead to decreased insulin production, β -cell dysfunction and metabolic abnormalities [13].

Membrane Potential, Ion Channels and Oxidative Stress

Oxidative stress is an essential method by which diabetes may impair the electrical activity of cells. Chronic hyperglycaemia may augment the creation of reactive oxygen species and mitochondrial dysfunction may further affect ATP synthesis and the redox homeostasis of the cells[14].

Since ATP is directly involved in control of K_{ATP} channels, alterations in mitochondrial metabolism may influence membrane potential. Oxidation damage of membrane proteins, composition alterations of lipids and alteration of intracellular calcium may also alter ion-channel function[15].

Association of membrane potential with antioxidant related measures in diabetes has been observed, supporting the hypothesis that electrical properties of membrane may represent wider metabolic and oxidative disturbances[16].

Ion Channels and Glucose Homeostasis Outside the β -Cell

Ion channels are not exclusive to pancreatic β -cells. They play a significant role in vascular smooth muscle, endothelial cells, neurones, skeletal muscle and other tissues involved in glucose metabolism[17].

Potassium channels in vascular tissues are involved in membrane polarisation and control calcium entry, contraction and relaxation of the vessels. Therefore, alterations in ion-channel function coupled to diabetes could be involved in the vascular dysfunction.

In the neurological system, potassium and calcium channels are responsible for neurone excitability, neurotransmitter release and signal transmission. Changes in these systems can contribute to the development of diabetic neuropathy complications[18].

Ion channels may also indirectly affect insulin sensitivity by impacts on vascular perfusion, neural control and cellular metabolism. Thus, membrane potential should be regarded as a property of an integrated physiological network and not an individual property of pancreatic β -cells [19].

Membrane Potential as a Biomarker Candidate

Glucose and glycated haemoglobin are the traditional laboratory assessment of diabetes and both are significant in giving information on glycaemic state. However, these assessments are, for the most part, metabolic outputs instead of the physiological mechanisms responsible for altered glucose regulation[20].

The membrane potential is of interest in that it represents the functional status of ion channels and cell membranes. The membrane potential was presented as a possible laboratory biomarker in type 2 diabetes, relating it to β -cell activity, ion channels, oxidative stress and glycaemic control[21].

Moreover, although further research is needed before membrane-potential measurements can be routinely integrated into clinical diagnosis, the investigation of membrane electrical characteristics may provide significant information concerning cellular dysfunction associated with diabetes[22].

Therapeutic Implications

The study of ion channels has already made an impact on treatments for diabetes. For example, sulfonylureas function, in part, by inducing closure of pancreatic β -cell K_{ATP} channels, hence stimulating membrane depolarisation and insulin secretion. This pharmacological concept shows the therapeutic value of the association of ion channels and membrane potential[23].

Future strategies may aim at more targeted control of K_{ATP}, voltage-gated calcium, potassium and other ion channels, while retaining normal β -cell electrical activity. Strategies to preserve mitochondrial function to reduce oxidative stress and preserve intracellular calcium homeostasis may also indirectly safeguard ion-channel function. However, since ion channels are expressed in the whole body, therapeutic modulation should be selective enough to minimise adverse effects on cardiac, vascular, neuronal and other tissues[24,25].

Membrane Potential as a Biomarker Candidate

Glucose and glycated haemoglobin are the traditional laboratory assessment of diabetes and both are significant in giving information on glycaemic state. However, these assessments are, for the most part, metabolic outputs instead of the physiological mechanisms responsible for altered glucose regulation[20].

The membrane potential is of interest in that it represents the functional status of ion channels and cell membranes. The membrane potential was presented as a possible laboratory biomarker in type 2 diabetes, relating it to β -cell activity, ion channels, oxidative stress and glycaemic control[21].

Moreover, although further research is needed before membrane-potential measurements can be routinely integrated into clinical diagnosis, the investigation of membrane electrical characteristics may provide significant information concerning cellular dysfunction associated with diabetes[22].

Therapeutic Implications

The study of ion channels has already made an impact on treatments for diabetes. For example, sulfonylureas function, in part, by inducing closure of pancreatic β -cell K_{ATP} channels, hence stimulating membrane depolarisation and insulin secretion. This pharmacological concept shows the therapeutic value of the association of ion channels and membrane potential[23].

Future strategies may aim at more targeted control of K_{ATP} , voltage-gated calcium, potassium and other ion channels, while retaining normal β -cell electrical activity. Strategies to preserve mitochondrial function to reduce oxidative stress and preserve intracellular calcium homeostasis may also indirectly safeguard ion-channel function.

However, since ion channels are expressed in the whole body, therapeutic modulation should be selective enough to minimize adverse effects on cardiac, vascular, neuronal and other tissues[24,25].

Conclusion

Membrane potential is an important physiological connection between glucose metabolism and insulin production. In pancreatic β -cells, K_{ATP} channels couple changes in cellular ATP synthesis to changes in membrane potential, while voltage-dependent calcium channels couple depolarisation to intracellular Ca^{2+} signals that activate insulin release. Then other potassium and calcium channels control repolarisation of the membrane, calcium dynamics, and the pattern of electrical activity.

In diabetes mellitus, prolonged hyperglycaemia, oxidative stress, lipotoxicity, mitochondrial dysfunction and altered calcium homeostasis might destabilise the highly regulated system. The ensuing ion-channel and membrane-potential anomalies might contribute to impaired insulin production and eventual β -cell failure. Because identical ion-channel processes exist in vascular and neural tissues, their malfunction may also play a role in the diabetic consequences. Understanding the biology of membrane potential and ion-channels thus provides a crucial biological framework for understanding glucose homeostasis and diabetes mellitus. Additional research on the membrane potential, activity of potassium and calcium channels may aid in the identification of novel biomarkers and conventional therapies for the prevention and treatment of diabetes.

References

1. Nnodim, J., Maduiké, C.V. and Nnodim, A.P. (2026). Membrane potential as a potential laboratory biomarker in type 2 diabetes mellitus: biological basis and clinical relevance. *Nexus Global Research Journal of Medical Science*, 2(9).
2. Nnodim, J. and Ozoemena, C.C. (2026). Calcium signaling in diabetes: role of ion channels and transporters. *Endocrine System and Diabetes*, 4(1).
3. Nwanyaka, H.I. and Nnodim, J. (2025). Correlationship of membrane potential, malondialdehyde and ascorbic acid in people living with diabetics in Owerri. *SAR Journal of Surgery*, 6(1), pp.1–3.
4. Ashcroft, F.M. and Rorsman, P. (2013). K_{ATP} channels and islet hormone secretion: new insights and controversies. *Nature Reviews Endocrinology*, 9, pp.660–669.
5. Rorsman, P. and Ashcroft, F.M. (2018). Pancreatic β -cell electrical activity and insulin secretion: of mice and men. *Physiological Reviews*, 98(1), pp.117–214.
6. Nnodim, J., Osuala, P.C. and Nwaokoro, J.C. (2022). Perspective of channelopathy in diabetes mellitus. *Biomedical Journal of Scientific & Technical Research*, 46(5).
7. Nnodim, J.K. and Ugwuibe, N.J. (2022). Disorders associated with ion channels. *Journal of Clinical and Medical Reviews*, 1(2).
8. Ashcroft, F.M. (2007). ATP-sensitive K^{+} channels and disease: from molecule to malady. *American Journal of Physiology-Endocrinology and Metabolism*, 293(4).
9. Gloyn, A.L., Siddiqui, J. and Ellard, S. (2006). Mutations in the genes encoding the pancreatic beta-cell K_{ATP} channel subunits Kir6.2 (KCNJ11) and SUR1 (ABCC8) in diabetes mellitus and hyperinsulinism. *Human Mutation*, 27, pp.558–567.
10. Bennett, K., James, C. and Hussain, K. (2010). Pancreatic β -cell K_{ATP} channels: hypoglycaemia and hyperglycaemia. *Reviews in Endocrine and Metabolic Disorders*, 11, pp.157–163.
11. Nnodim, J.K. and Opara, A.U. (2012). The plasma vitamin C and E status in Type II diabetes with malaria in Owerri, Nigeria. *Pakistan Journal of Nutrition*, 11(1), pp.62–63.

12. Nnodim, J.K., Nnanemere, I., Ihim, A., Okorie, H. and Udujih, H.I. (2012). Plasma glucose level and antioxidant vitamins C and E in an alloxan-induced diabetic mellitus rats. *Indian Journal of Fundamental and Applied Life Sciences*, 2(1), pp.163–169.
13. Nnodim, J.K., Meludu, S.C., Dioka, C.E., Onah, C., Chilaka, U.J. and Obi, P.C. (2014). Altered membrane potential and electrolyte in sickle cell anemia. *Journal of Krishna Institute of Medical Sciences University*, 3(1), pp.70–73.
14. Remedi, M.S. and Nichols, C.G. (2022). ATP-sensitive potassium channels in hyperinsulinism and type 2 diabetes: inconvenient paradox or new paradigm? *Diabetes*, 71(3), pp.367–375.
15. Nnodim, J.K., Nwaokoro, J.C., Edward, U. and Oly-Alawuba, N. (2020). Evaluation of selectins, membrane potential and antioxidant vitamins in diabetic patients attending General Hospital Owerri. *EC Diabetes and Metabolic Research*, 4(10), pp.26–30.
16. Nnodim, J.K., Njoku, C.J., Nwaokoro, J.C. and Ihionu, C.A. (2019). Evaluation of selenium, calcium and membrane potential among breast cancer patients in Owerri. *Global Journal of Reproductive Medicine*, 7(2), pp.34–36.
17. McTaggart, J.S., Clark, R.H. and Ashcroft, F.M. (2010). The role of the K_{ATP} channel in glucose homeostasis in health and disease: more than meets the islet. *The Journal of Physiology*, 588(17), pp.3201–3209.
18. Seino, S. and Miki, T. (2003). Physiological and pathophysiological roles of ATP-sensitive K⁺ channels. *Progress in Biophysics and Molecular Biology*, 81, pp.133–176.
19. Ashcroft, F.M. (2007). ATP-sensitive potassium channelopathies: focus on insulin secretion. *Journal of Clinical Investigation*, 117, pp.2047–2058.
20. Satin, L.S., Butler, P.C., Ha, J. and Sherman, A.S. (2015). Pulsatile insulin secretion, impaired glucose tolerance and type 2 diabetes. *Molecular Aspects of Medicine*, 42, pp.61–77.
21. Zhou, Y., Sun, P., Wang, T., Chen, K., Zhu, W. and Wang, H. (2015). Inhibition of calcium influx reduces dysfunction and apoptosis in lipotoxic pancreatic β -cells via regulation of endoplasmic reticulum stress. *PLoS ONE*, 10, e0132411.
22. Ravier, M.A. and Rutter, G.A. (2014). Calcium signalling in pancreatic β -cells in health and in Type 2 diabetes. *Cell Calcium*, 56(5), pp.340–361.
23. Thévenod, F. (2011). Calcium channels and transporters in the pathogenesis of diabetes mellitus. *Cell Calcium*, 50, pp.111–122.
24. Braun, M., Ramracheya, R., Bengtsson, M., et al. (2008). Voltage-gated ion channels in human pancreatic beta-cells: electrophysiological characterization and role in insulin secretion. *Diabetes*, 57(6), pp.1618–1628.